

The Maine CDC/DHHS Cardiovascular Health Program (MCVHP) works to reduce death and disability due to heart disease and stroke among Maine residents. We have developed this fact sheet to help you understand the burden of cardiovascular disease in Maine and your community by presenting key data.

KEY FINDINGS

- Compared to all Maine residents, persons in Western District are more likely to be hospitalized for major CVD, coronary heart disease, and heart attack.
- Compared all Maine adults, a lower percent of adults in Western District said their blood cholesterol was checked within the previous five years.
- Compared all Maine adults, a lower percent of adults in Western District with high blood pressure also said they were taking prescription drugs for blood pressure control.

Major cardiovascular diseases (heart disease and stroke) are the leading cause of death among men and women in the United States and in Maine, accounting for roughly 33 percent and 28 percent of all deaths, respectively, in 2005¹. Some risk factors that you can change for cardiovascular disease include lack of physical activity, poor nutrition, tobacco use, high blood pressure, high blood cholesterol, overweight, and diabetes. Data related to high blood pressure and high blood cholesterol are included in this fact sheet. In addition, we present data on knowledge about warning signs and symptoms for heart attack and stroke, as well as death and hospitalization rates for selected cardiovascular diseases.

Prevalence of Cholesterol Indicators Among Adults – Western District, 2005

High levels of cholesterol in the blood can lead to blockage of the arteries and cause diseases such as coronary heart disease, heart attack and stroke.

Cholesterol Screening


In 2005, about 72 percent of adults in Western District said their blood cholesterol was checked within the previous five years. This is lower than the percent of Maine adults who had their cholesterol checked. More women than men had their cholesterol checked. Adults with college education or high household income are more likely to have had their cholesterol checked.

High Cholesterol

In 2005, about 43 percent of adults in Western District had high cholesterol levels. This is higher but comparable to the percent of Maine adults who have high cholesterol. Fewer women have high cholesterol than men. Adults with low household income or less than college education are more likely to have high cholesterol levels.

Cholesterol Control

In 2007, nearly 96 percent of adults in Western District with high blood cholesterol said they were following their doctor's advice to take prescribed medication to control their cholesterol levels. This is higher than the percent of Maine adults with high cholesterol who said they were taking the same action following physician advice. There were no differences between men and women. Adults with household income of less than \$25,000 or less than a high school education are more likely to take medications to control blood cholesterol levels.



Prevalence of Blood Pressure Indicators Among Adults – Western District, 2005

High blood pressure is known to increase the risk of heart disease and stroke.

Blood Pressure Screening


According to data from the 2000 Maine BRFSS, most adults in Maine check their blood pressure at least once every two years with no apparent evidence of disparities across subpopulations.

High Blood Pressure

In 2005, nearly 25 percent of adults in Western District had high blood pressure. This is comparable to the percent of Maine adults who have high blood pressure. Fewer women than men have high blood pressure. Adults with low household income or less than college education are more likely to have high blood pressure.

Blood Pressure Control

In 2005, more than 74 percent of adults in Western District with high blood pressure also said they were taking prescribed medications to control their blood pressure. This is comparable to the percent of Maine hypertensive adults who take medications to control their blood pressure. More women than men take medications for blood pressure control. Adults with low household income or less than college education are more likely to take medications to control their blood pressure.



2005 Maine Behavioral Risk Factor Surveillance System

Prevalence of Knowledge About Heart Attack and Stroke Among Adults – Western District, 2005


Early recognition of the warning signs and symptoms of heart attack or stroke by patients and by bystanders are vital to timely access to emergency care and receiving of lifesaving treatment.^{2,3}

Warning Signs and Symptoms for Heart Attack

In 2005, more than 14 percent of adults in Western District were able to correctly identify the major signs and symptoms of heart attack. This is similar to the percent of Maine adults who were able to identify the same set of signs and symptoms. Compared to men, more women are able to identify the signs and symptoms of heart attack. Generally, adults between the ages of 35 and 64 years are more likely to correctly identify the signs and symptoms of heart attack compared to those 18 to 34 or 65 and older.

Warning Signs and Symptoms for Stroke

In 2005, nearly 21 percent of adults in Western District were able to correctly identify the major signs and symptoms of stroke. This is similar to the percent of Maine adults who identified the same set of warning signs and symptoms. Women appear more knowledgeable about the warning signs and symptoms of stroke. Adults between the ages of 35 and 64 years are more likely to correctly identify the signs and symptoms of stroke compared to those younger than 35 or 65 and older.



2005 Maine Behavioral Risk Factor Surveillance System

Hospitalizations Due to Major Cardiovascular Disease (CVD), Coronary Heart Disease (CHD), Heart Attack, and Stroke – Western District, 2005

Hospitalization rates for CVD are a measure of disease burden. Hospitalizations can be minimized with early detection and appropriate management of CVD.

Major Cardiovascular Diseases

In 2005, more than 3,380 residents of Western District were hospitalized for major CVD. The unadjusted hospitalization rate was about 1740 per 100,000. The age-adjusted hospitalization rate is lower than the statewide rate.

Coronary Heart Disease


In 2005, more than 1,240 residents of Western District were hospitalized for coronary heart disease. The unadjusted hospitalization rate was about 639 per 100,000. The age-adjusted hospitalization rate is lower than the statewide rate.

Heart Attack

In 2005, more than 520 residents of Western District were hospitalized for acute heart attack. The unadjusted hospitalization rate was about 270 per 100,000. The age-adjusted hospitalization rate is lower than the statewide rate.

Stroke

In 2005, nearly 510 residents of Western District were hospitalized for stroke. The unadjusted hospitalization rate was about 261 per 100,000. The age-adjusted hospitalization rate is similar to the statewide average.



Maine Health Data Organization

Deaths Due to Major Cardiovascular Disease (CVD), Coronary Heart Disease (CHD), Heart Attack, and Stroke – Western District, 2005

CVD death rates are another measure of disease burden. Death from CVD can be minimized through primary prevention, early detection, appropriate management of CVD, and timely access to quality care for acute events.

Major Cardiovascular Diseases

590 residents of Western District died of major CVD in 2005. This corresponds to an unadjusted death rate of about 304 per 100,000. The age-adjusted death rate is comparable to the statewide average.

Coronary Heart Disease


About 290 residents of Western District died of coronary heart disease in 2005, representing an unadjusted death rate of 148 per 100,000. The age-adjusted death rate is similar to the statewide average.

Heart Attack

More than 120 residents of Western District died of acute heart attack in 2005, representing an unadjusted death rate of about 62 per 100,000. The age-adjusted death rate is comparable to the statewide average.

Stroke

110 residents of Western District died of stroke in 2005, representing an unadjusted death rate of about 57 per 100,000. The age-adjusted death rate is comparable to the statewide average.



Maine Office of Data, Research and Vital Records

The Cost of Cardiovascular Disease

In 2005, national indirect cost of cardiovascular disease due to mortality was estimated at \$117 billion.⁴ Based on national figures, the indirect cost of cardiovascular disease mortality in Maine for 2005 was estimated at \$501 million.⁴ This figure includes the cost of lost productivity as a result of premature deaths.

Prevalence of cholesterol indicators among adults – Western District, 2005

	Checked cholesterol within last 5 years	High cholesterol
Androscoggin	71.0	39.1
Franklin	76.8	51.4
Oxford	70.7	44.9
Western District	71.8	42.9
Maine	78.7	36.2

Prevalence of blood pressure indicators among adults – Western District, 2005

	High blood pressure	Take medication for blood pressure
Androscoggin	25.4	71.6
Franklin	28.8	66.0
Oxford	21.8	86.8
Western District	25.0	74.3
Maine	25.8	76.5

Prevalence of knowledge about heart attack and stroke among adults – Western District, 2005

	Identify 5 major heart attack signs and decoy	Identify 5 major stroke signs and decoy
Androscoggin	13.2	19.5
Franklin	15.9	21.1
Oxford	17.4	17.6
Western District	14.4	20.7
Maine	13.4	36.2

Age-adjusted hospital discharge rates due to major CVD, coronary heart disease (CHD), heart attack, and stroke – Western District, 2005

	Major CVD	CHD	Heart Attack	Stroke
Androscoggin	1388.5	488.6	203.9	205.9
Franklin	1808.7	778.6	379.5	210.7
Oxford	1518.2	533.9	209.6	258.8
Western District	1486.5	543.1	229.5	222.0
Maine	1560.3	620.4	292.4	207.0

Age-adjusted death rates due to major CVD, coronary heart disease (CHD), heart attack, and stroke – Western District, 2005

	Major CVD	CHD	Heart Attack	Stroke
Androscoggin	234.8	116.3	41.1	41.9
Franklin	265.2	108.9	68.0	*
Oxford	278.5	145.8	64.1	51.2
Western District	251.4	123.9	51.4	46.5
Maine	242.0	124.1	59.4	42.9

*Rates are suppressed when the number of events is <20

- 1 H.C. Kung, Ph.D., D.L. Hoyert, Ph.D., J.X., M.D., S.L. Murphy, B.S. Deaths: Final Data for 2005. National vital statistics reports. Hyattsville, MD. National Center for Health Statistics, 2008.
- 2 W. Hacke, G. Donnan, C. Fieschi, et al. Association of outcome with early stroke treatment: pooled analysis of ATLANTIS, ECASS, and NINDS rt-PA stroke trials. Lancet 2004; 363: 768-774.
- 3 Fang, N. Keenan, S. Dai, C. Denny. Disparities in Adult Awareness of Heart Attack Warning Signs and Symptoms – 14 States, 2005. MMWR 2008; 57: 175-179.
- 4 American Heart Association: Heart Disease and Stroke Statistics 2005 Update. Dallas, TX: American Heart Association; 2008. Available at: <http://www.americanheart.org/downloadable/heart/1105390918119HDSStats2005Update.pdf>.

Technical Notes: Age-adjusted rates are adjusted to the year 2000 United States standard population. All subpopulation statements appearing in district level fact sheets are based on Maine data and do not reflect actual disparities, if any, within the district.



*Maine Center for Disease
Control and Prevention*
An Office of the
Department of Health and Human Services

John E. Baldacci, Governor

Brenda M. Harvey, Commissioner

Printed under appropriation #013-10A-2626-032.

The Department of Health and Human Services (DHHS) does not discriminate on the basis of disability, race, color, creed, gender, age, sexual orientation, or national origin, in admission to, access to or operation of its programs, services, activities or its hiring or employment practices. This notice is provided as required by Title II of the Americans with Disabilities Act of 1990 and in accordance with the Civil Rights Acts of 1964 as amended, Section 504 of the Rehabilitation Act of 1973 as amended, the Age Discrimination Act of 1975, Title IX of the Education Amendments of 1972 and the Maine Human Rights Act. Questions, concerns, complaints, or requests for additional information regarding civil rights may be forwarded to the DHHS' ADA Compliance/EEO Coordinator, State House Station #11, Augusta, Maine 04333, 207-287-4289 (V) or 207-287-3488 (V), TTY: 800-606-0215. Individuals who need auxiliary aids for effective communication in programs and services of DHHS are invited to make their needs and preferences known to the ADA Compliance/EEO Coordinator. This notice is available in alternate formats, upon request.